

296#B

TRIACHLOR M

Emulsifiable Concentrate

5 ltrs

TPRI Registration No:

Pre-Emergence Herbicide for control of many important annual grasses and broadleaf weeds in sugar-cane and green beans

COMMERCIAL AND AGRICULTURAL CLASS / SEHEHU YA BIASHARA NA KILIMO

Contains 960g of Metolachlor per litre
Shake well before use

Lita moja ina gramu 960g za Metolachlor
Tikisa dawa vizuri kabla ya kutumia

Kiuagugu kinachoangamiza aina nyingi za magugu yenye majani mapana na nyasi kabla hayaajaota, katika zao la miwa na maharage



Supplied by Tradebase Ltd
Sussex House, Guildford
GU3 3BH England

TRIACHEM (T) LTD
Esso Road
P.O. Box 2750
Arusha
Tel/Fax: +255 27 250 2446

Date of Manufacture: NOV.2006
Batch No: 20061102
Expiry Date: NOV.2009

**ALWAYS READ THE LABEL BEFORE USING
SOMA KIBANDIKO CHA MAEALEZO KABLA YA KUTUMIA DA WA HI
KEEP LOCKED UP AND OUT OF REACH OF CHILDREN
WEKA MBALI NA WATOTO**



APPLICATION RATE

GREEN BEANS/ SUGARCANE	0.75 - 2.0LTS/HA applied in 120-600ls of water.
---------------------------	---

DIRECTIONS FOR USE

- APPLY to soil before, at, or immediately after planting and before crop/weeds emerge. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4 cm of soil.
- Use high rates on heavy soils or where heavy grass infestation is expected.
- Use low rates on soils where sand content is in 50-70% range.
- Use sufficient agitation during mixing and application to maintain a uniform emulsion.
- DO NOT apply to waterlogged soils
- DO NOT apply when raining heavily.
- DO NOT irrigate to point of run-off for at least 2 days after application.

WEEDS CONTROLLED

Digitaria filiformis, ischaemum, horizontalis, sanguinalis
Panicum dichotomiflorum, laevifolium.
Setaria faberi, glauca, pallidifusca, verticillata, viridis.
Brachiaria eruciformis; Chloris virgata; Echinochloa crus-galli.
Gallinsoga parviflora; Nicandra physaloides; Urochloa panicoides.

Note: Limited control at the higher rate will also be given against *Amaranthus* Sorghum and *Cyperus* species.

SAFETY PRECAUTIONS

- Keep locked up and out of reach of children.
- Keep only in original container away from food, drink and animal feed.
- When handling do not eat, drink or smoke.

PRECAUTIONS WHEN MIXING

- Wear rubber gloves and eye protection.
- If contacts skin or eyes wash immediately with plenty of water.
- If product spills on clothes change and wash them.

PRECAUTIONS WHEN SPRAYING

- Wear boots long trousers, long sleeved shirt and eye protection.
- Avoid contact with spray.
- Keep away from ponds, waterways, wells and ditches.

PRECAUTIONS AFTER SPRAYING

- Wash yourself and your clothes.
- Do not re-use this container. Triple rinse, destroy by cutting and then bury.

ACCIDENTAL RELEASE MEASURES

- Wear protective clothing when dealing with spillages.
- Prevent entry into drains and water-courses by containing spillage with sand/soil.
- Absorb spill chemical with dry sand/earth
- Remove contaminated sand/soil and bury in uncropped land away from water supplies.

FIRST AID

If product is swallowed drink plenty of water and visit your doctor immediately taking this label with you.

MEDICAL TREATMENT

Metolachlor is slightly toxic and has no specific antidote. Treat symptomatically and administer emesis or stomach lavage as appropriate. 5mg/kg of activated charcoal suspension (50g/400ml water) can be given to absorb the remaining product.

MATUMIZI:

MAHARAGE/MIWA	Tumia 0.75 - 2.0L/ta/Hektari katika lita 120-600 za maji.
---------------	---

JINSI YA KUTUMIA:

- Nyunyiza dawa katika udongo kabla, mara baada ya kupanda na kabla mimea/magugu kooa. Ni vyema mvua ikinyesha/umwagilaji kufanyika siku 10 baada ya kupanda ili kutowanisha udongo wa juu 3-4cm.
- Tumia kiwango cha juu cha dawa kwenye udongo wa mfinyanzi, pia Schema ambao zinategemea kuota magegu mengi.
- Tumia kiwango cha chini cha dawa kwenye udongo wenye mchanga kati ya asilimia 50 - 70
- Kuroga dawa vizuri wakati wa kuchanganya, ili kupata uiano mzuri wa dawa na maji.
- USINYUNYZE DAWA katika udongo uliosimamisha maji
- USINYUNYZE DAWA wakati mvua inayesha.
- USINYESHE maji mengi sana kabla ya siku 2 baada ya kunyunyiza dawa.

MAGUGU YANAYOTA NGAMIZWA:

Digitaria filiformis, ischaemum, horizontalis, sanguinalis
Panicum dichotomiflorum, laevifolium.
Setaria faberi, glauca, pallidifusca, verticillata, viridis.
Brachiaria eruciformis; Chloris virgata; Echinochloa crus-galli.
Gallinsoga parviflora; Nicandra physaloides; Urochloa panicoides.

Dokezo: Utumizi kwanza cha juu cha dawa hii, kinaweka kuhitaji magegu aina ya Mchicha pori, Mtama pori, pamoja na Ndogo.

USALAMA KWA WANAYUNYUNYZA:

- Pungia dawa mahali mbali na watoto.
- Hifadhi dawa kwenye vyombo vyake atili mahali mbali na vyakula, vinywaji pamoja na wanyama.
- Usile, usinywe wala kuvuta sigara wakati unaponyunyiza dawa.

TABADHA RI WAKATI WA KUCHANGANYA:

- Vaa glava mikononi, na ngao ya uso.
- Ukimwagikiwa na dawa kwenye ngozi, au machosi nawa mara moja kwa kutumia maji mengi.
- Ukimwagikiwa na dawa ngooni, bafilisha ngao kisha uzifite.

TABADHA RI WAKATI WA KUNYUNYIZA:

- Vaa gam boots, suruali safeti, shati la mikono mirefu na ngao ya uso
- Epuka kugusana na dawa.
- Ni sumu kwa viembe wa majini, usitanyize dawa kwenye mito au mitaro ya maji.

TABADHA RI BAADA YA KUNYUNYIZA:

- Oga mwili mzima, kisha zifae ngao zako.
- Usitumie tena chombo kilichoisha dawa, hariba chombo kwa kukitoboa toba na ukifukie.

HATUA ZA KUFUATA DAWA IKIM WAGIKA KWA BAHATI MBAYA:

- Vaa vifaa vya kinga wakati unapohughulika dawa iliyomwagika.
- Epuka dawa iliyomwagika isingie kwenye mitaro na vyanzo vya maji kwa kuchanganya na mchanga au udongo.
- Tumia viloyota maji kama mchanga au udongo mkavu kwa kufyonza dawa iliyomwagika.
- Ondoa mchanga/udongo baada ya kufyonza dawa iliyomwagika na ufukie schemu itiyu na maza mbali na vyanzo vya maji.

HUUMA YA KWANZA:

Iwapo dawa itamewa inashauriwa kwenya maji mengi na mgonjwa aplekwe kwa dakitari mara moja pamoja na kibandiko cha dawa.

HUUMA KWA DAKITARI:

Metolachlor siyo dawa yenye sumu kali, na haina makata maalum. Mgonjwa athiwe tu kulingana na matatizo yatakavyojikwa, na mgonjwa atapibwe au apewe dawa ya kuhitisha 5mg/kg za activated charcoal suspension (50g/400ml za maji) ili kufyonza mabaki ya dawa tumboni.