

Lebaycid[®] EC 500

Fenthion 500g/l

Lebaycid[®] is a broad spectrum insecticide with good persistence and depth action for use in coffee, vegetables, fruit crops, and other crops for control of sucking and chewing pests.



Aphids



Antestia



Berry Borer



Leaf miner

Main Features

- Acts as a contact and stomach poison.
- Displays good initial activity.
- Has long persistence and good depth action.
- Well tolerated by plants when used at recommended rates.
- Soft on beneficials.



Thrips

Compatibility

Lebaycid[®] is compatible with Bayer insecticides and fungicides e.g Antracol, Baycor and Bayfolan. Spray mixture should be applied immediately.

Application

Apply as a full cover spray in recommended volume of water ensuring even spray coverage of all plant surface. Water volume : 1,000 litres per hectare.

Directions for use:

Crop	Pest	Rate/ha	Rate/20lt water	Timing
Coffee	Leafminer, mealybug, Berry moth, Berry borer	2.25lt	45ml	Apply as soon as the pests are seen and repeat as required
	Thrips	0.75lt	15ml	
	Antestia, Lace bug, Stinging Caterpillar, Jelly grub, Leaf Skeletoner, Tailed caterpillar	0.9-1.25lt	20-25ml	
Vegetable	Aphids, Thrips, Leafminers, Fruitfly, Melon fly Caterpillars	0.9-1.25lt	20-25ml	Apply as soon as the pests are seen and repeat at 14-21 day interval as required
Tobacco	Aphids, Thrips,	0.9-1.25lt	20-25ml	Apply as soon as the pests are seen and repeat 21 day interval as required
Citrus, Mangoes, Guavas, Plums, Peaches, Apples, Pears	Fruitfly, Aphids, Caterpillars, False codling moth,	0.9-1.25lt	20-25ml	Apply as soon as the pests are seen and repeat 21-28 day interval as required
	Thrips	1.25lt	25ml	
Rice	Stem borers	0.9-1.25lt	20-25ml	Apply as soon as the pests are seen and repeat 21 day interval as required
Sugarcane	Sugarcane borers	0.9-1.5lt	20-30ml	Apply as soon as the pests are seen and repeat as required
Grapes	Fruit fly, Caterpillar	0.9-1.25lt	20-25ml	Apply as soon as the pests are seen and repeat 21 day interval as required
Maize	Stem borers	1.5-2.0lt	30-40ml	Apply at 3-4weeks after germination and repeat once 21 days later if required
All crops	Fruit fly (bait treatment)	0.75lt	15ml	Mix with Buminal bait and apply as a coarse bait spray. Repeat after 21 days if required

Pre-harvest Interval

Allow 14 days between the final application of Lebaycid® and harvest of any one food crop.