

NURELLE* D 50/500EC

Mode of Action: Contact
Stomach Action
Vapour Action

DIRECTIONS FOR USE

Nurelle D 50/500EC can be applied through any medium or high volume conventional 'ground' sprayer (200-1000l/ha) or through 'aerial' application equipment (minimum 50lt/ha). Being a contact insecticide, good coverage is essential especially for under leaf sucking insects and where crop canopy is dense.

TIMING OF APPLICATION

Apply Nurelle D 50/500EC at first sign of pest attack. Spray during cool hours of the day (early morning and late afternoon). Do not spray if rain is expected within an hour.

RATES OF APPLICATION

CROP	PEST	RATE/HA	RATE/20LT	REMARKS
COTTON	American bollworm Spiny bollwor Red bollworms Army worm Cotton leafworm	500ml	20ml	Control will be improved by early application.
	Cotton aphid Whitefly Thrips Cotton jassid Cotton stainers Capsids	800ml	30ml	Long term control is improved by repeat application at 12-14 days interval.
	Red spider mites	800ml	30ml	Suppression will be effective with repeat spraying. It prevents flare up of mites.
Vegetables Beans	Aphids, ants, whitefly, cabbage moth, bollworms, leaf eater, caterpillars, bean fly.	400-800ml	8-16ml	Spray according to pest threshold and repeat spray at 12-14 days according to infestation.
Wheat	Aphids Bollworms	400-800ml	16ml	Higher rates for aphids.

SPRAY VOLUME

Spray in 800-1000litres of water per hectare.

Pre harvest period (PHI)

Vegetables 7-14 days

Re-entry period

6 hours

PACKS AVAILABLE

100ml, 500ml 1lt, 20Lt, 200Lt

ALWAYS READ THE LABEL BEFORE USE

*Trademark of Dow AgroSciences

Marketed By:

Lachlan Kenya Limited
P.O. Box 49470 00100
Nairobi-Kenya
Tel: 254-20-823105/419/106
Fax: 254-20-823093
e-mail: lachlan@griculture.co.ke
website: www.griculture.co.ke

Distributed By:

 **Lachlan@griculture**

 **Dow AgroSciences**