

567932 KE/12E



sequestrene® 138 Fe 100 SG



1 kg

syngenta



For correction of iron deficiency in all kinds of crops and ornamentals in alkaline and calcareous soils (pH 2.5-11)

Water soluble granules

Micro-nutrient

GUARANTEE: EDDHA Na Fe (Iron (Fe) 6%)

Shelf Life

At least 2 years from date of manufacture in original sealed containers.

Storage

Store away from sun and damp. Keep out of reach of children and animals.

Avoid storage above +35°C.

® Registered Trademark of a Syngenta Group Company

© 2001 Syngenta Crop Protection AG

Manufactured by:

Syngenta Crop Protection AG, Basle, Switzerland

Distributor:

Syngenta East Africa Ltd

P. O. Box 30393

00100 - NAIROBI

Tel: 254-02-532750/532983

Fax: 254-02-532753/535456



Mode of Action

Sequestrene® is taken up by the roots and leaves, but the root uptake gives more persistent effect than can be achieved with foliar treatments.

Directions for Use

Sequestrene® can be applied to the roots or to the foliage of all kinds of annual and perennial crops and ornamentals. The treatment can either be curative or preventive. It corrects iron deficiencies and increases the quantity and quality of harvest. Sequestrene® is preferably applied around planting time or any time

Sequestrene® Fe 138 for curative use

Crop Mimea	Sequestrene® Kiasi cha kutumia	Remarks Maeleo
Vegetables Mboga za and / na	g/100 lt. 200-300	Foliar spray Kunyuyizia majani aina mbali mbali
Ornamentals Maua	kg/ha. 2.0-5.0	Soil application Kuweka kwa mchanga
Tree crops - Citrus, vines Matunda ya aina mbali mbali	g/tree for each cm of tree trunk diameter	Do not exceed 0.5kg/tree. Reduce rate by 30-40% in cases of less severe chlorosis
Field Crops- wheat, maize Mimea mengine	0.5 2.0-5.0	Usizidi nusu kilo kwa mti mmoja. Punguza kiasi kwa 30-40% kama upungu si mwingi. Foliar spray with sufficient water for full coverage. Kunyuyizia majani na maji ya kutosha Soil application Kuweka kwa mchanga

when plants are actively growing.

Sequestrene® as fertilizer/growth stimulant.

Apply 20 kg/ha broadcast or hand sprayed or dry soil incorporated or not. Do not exceed this rate because non-absorbed Sequestrene® 138 Fe is washed out.

Timing

Preventive Use - Apply early in the season either at planting or any time during active growth of the plants.
Curative use - Ascertain that the chlorosis is a result of iron deficiency. Sequestrene® can be applied curatively any time so long as the environment is conducive to active growth.

Operator Safety

Do not eat, drink or smoke during work. Avoid breathing spray mist and spray dust. Wash hands and face thoroughly before eating, smoking or drinking. Wear protective clothing, i.e. overalls, gloves, faceshield, cap, etc, when working with the product. Change and wash clothing before re-use.

User's Risk

The manufacturer warrants the quality of their product so long as it is acquired in the original sealed container. Many factors such as type of soil, weather, type of plant, method of application, may affect the performance of Sequestrene®. The user bears the risk for damage resulting from such factors which are beyond the manufacturers control. The manufacturer declines all liability resulting from improper storage or use of the product.


® Registered Trademark of a Syngenta Group Company

© 2001 Syngenta Crop Protection AG

567932 KE/12E



sequestrene® 138 Fe 100 SG



Mbolea ya kupungua ukosekani wa chakula ya aina ya chuma kwenvye maua na mimiea mingine (pH 2.5-11).

Viduchu viowevu kwa maji

DHAMANA: EDDHA Na Fe (Iron (Fe) 6%)

Maisha rafuni

Si chini ya miaka miwili kuanzia tarehe ya kutengenezwa ikihifadhiwa ndani ya mfuko au chombo chake asili.

Hifadhi mahali pakavu kivulini mbali na watoto, vyakula na mifugo. Joto katika sehemu ya kuhifadhi lisizidi +35°C.

® alama ya biashara ya kampuni ya Syngenta

© 2001 Syngenta Crop Protection AG

Watengenezaji:

Syngenta Crop Protection AG, Basle, Switzerland

Wauzaji:

Syngenta East Africa Ltd

P. O. Box 30393

00100 - NAIROBI

Tel: 254-02-532750/532983

Fax: 254-02-532753/535456



syngenta



Inavyofanya kazi

Sequestrene® huingia kwenye mizizi na majani ya mimea.

Inavyotumika

Sequestrene® inaweza kutumiwa kwa kuweka kwa mizizi au kwa kunyuyizia majani ya aina mbali mbali ya mimea au maua. Sequestrene® inaweza kutumiwa kuponya au kuzuia upungufu wa chuma katika mimea. Sequestrene® inapungua ukosekani wa chuma na huongea kiasi na ubora wa mavuno. Sequestrene® inapendekezwa kutumiwa wakati wa kupanda ama wakati mimea inakua haraka haraka.

Sequestrene® kama mbolea au ya kuchoecha kukua: Weka kilo 20 kwa hekta moja lioneze kote kote, ama kunyuyiza kwa mkono au kuchanganya na mchanga. Usizidi kipimo hiki kwa sababu Sequestrene® ambayo haitumiki na mimea hipotea bure.

Wakati wa kutumia

Kuzuu - tumia mapema wakati wa kupanda au wakati wovote mimea inakua haraka haraka. Kuponya - hakikisha shida inatokana na upungufu wa chuma. Sequestrene® Fe inaweza kutumiwa wakati wovote mimea inakua.

Usalama wa mfanyi kazi

Usinwe, usile wala kuvuta sigara unapofanya kazi. Epuka vumbi wa marashasha. Nawa mikono na uso kabla ya kula, kuvuta sigara au kunywa chocotte. Vaa nguo za kazi - yaani ovaroli, kofia, glau - hasa wakati wa kunyuyiza na kupaka Sequestrene®. Badilisha na usafishe nguo ulizova kabla ya kuzivaa tenu.

Dhamana

Watengenezaji wanathibitisha ubora wa Sequestrene® ukiipata ikiwa furusi yake hajafunguliwa. Hali nyinyi hasi zile zinazotegemea hali ya hewa, aina ya mchanga aina ya mimea na namna Sequestrene® inavyotumiwa huweza kuleta uharibifu au kutofaulu kwa Sequestrene®. Yoyote ya hayo yakiteka yatakuwa juu ya mtumiaji. Watengenezaji na wauzaji wanajikinga kwa lawama yoyote kwanzi hali hizi hazimo mikononi mwao.

® alama ya biashara ya kampuni ya Syngenta

© 2001 Syngenta Crop Protection AG, Basle, Switzerland